

Nuclear scattering of dark matter coupled to a new light scalar

Douglas P. Finkbeiner,^{1,*} Tracy R. Slatyer,^{2,†} and Neal Weiner^{3,‡}

¹*Harvard-Smithsonian Center for Astrophysics, 60 Garden St., Cambridge, MA 02138*

²*Physics Department, Harvard University, Cambridge, MA 02138*

³*Center for Cosmology and Particle Physics, Department of Physics, New York University, New York, NY 10003*

(Dated: October 17, 2008)

We consider the nuclear scattering cross section for the eXciting Dark Matter (XDM) model. In XDM, the Weakly Interacting Massive Particles (WIMPs) couple to the Standard Model only via an intermediate light scalar which mixes with the Higgs: this leads to a suppression in the nuclear scattering cross section relative to models in which the WIMPs couple to the Higgs directly. We estimate this suppression factor to be of order 10^{-5} . The elastic nuclear scattering cross section for XDM can also be computed directly: we perform this computation for XDM coupled to the Higgs sector of the Standard Model and find a spin-independent cross section in the order of 4×10^{-13} pb in the decoupling limit, which is not within the range of any near-term direct detection experiments. However, if the XDM dark sector is instead coupled to a two-Higgs-doublet model, the spin-independent nuclear scattering cross section can be enhanced by up to four orders of magnitude for large $\tan\beta$, which should be observable in the upcoming SuperCDMS and ton-scale xenon experiments.

PACS numbers: 95.35.+d

I. INTRODUCTION

Motivated by the apparent excess of e^+e^- annihilation in the Galactic center [1, 2], Finkbeiner & Weiner [3] proposed a model of eXciting Dark Matter (XDM) in which Weakly Interacting Massive Particles (WIMPs) collisionally excite and subsequently de-excite via e^+e^- emission. This model uses the kinetic energy of the WIMP dark matter to create e^+e^- pairs, in contrast with light dark matter models in which the pairs result from the mass energy of WIMP annihilation [e.g., 4], where the WIMP mass must be less than a few MeV [5, 6]. Because the XDM WIMP must have a weak-scale mass (~ 500 GeV) it retains many of the desirable properties of weak-scale WIMPs such as the thermal relic freeze-out abundance. Collisional excitation of weak-scale WIMPs was also discussed by Pospelov & Ritz [7].

The recent claim by DAMA/LIBRA of an annual modulation signal [8] conflicts strongly with other direct detection experiments [CDMS, XENON-10, ZEPLIN-II, CRESST], under the assumption of elastic nuclear scattering. Models in which the WIMP scatters *inelastically*, exciting an internal state of order 100 keV, appear to accommodate all available data [9]. The success of inelastic WIMP scattering in these two cases (albeit with different mass splittings) raises the question of whether a nuclear scattering cross section of 10^{-4} pb (required by Chang et al [9]) could arise naturally in the XDM model.

In XDM, WIMPs couple to the Standard Model via a new light scalar which mixes with the Higgs. Pre-

liminary results from ATIC [10] and PAMELA motivate the existence of just such a light particle [11]. However, the mixing angle between the light scalar and the Higgs suppresses the nuclear scattering cross section relative to generic models of dark matter where the Higgs couples directly to the WIMPs. We first estimate this suppression factor, then calculate the nuclear scattering cross section for the XDM dark sector coupled to the Standard Model or a two-Higgs-doublet model, and finally discuss the role of inelastic scattering and prospects for direct detection.

II. SUMMARY OF THE XDM MODEL

The defining feature of the XDM model is that the WIMP has an excited state which can be collisionally excited, and subsequently decay to produce e^+e^- pairs. The excited state could exist due to compositeness of the dark matter, or arise from an approximate symmetry of the theory.

For the excited state to be accessible in the Milky Way, and relevant for e^+e^- production, only a narrow kinematical range must be considered for the mass splitting, δ . For the decay to the ground state to be energetically capable of producing e^+e^- pairs, one must have $\delta > 1.022$ MeV. On the other hand, the kinetic energy available for a pair of 500 GeV WIMPs colliding each with velocities $v \sim 600$ km/s (roughly the escape velocity of the galaxy), is 2 MeV, setting an upper bound on δ .

To produce a sufficiently high number of positrons to explain the INTEGRAL signal, a large cross section is required [3, 7], comparable to the geometric cross section set by the characteristic momentum transfer. That is, $\sigma \sim (M_\chi \delta)^{-1}$ is of the correct size. Such a cross section can arise naturally [3], but requires the presence of a

*Electronic address: dfinkbeiner@cfa.harvard.edu

†Electronic address: tslatyer@fas.harvard.edu

‡Electronic address: neal.weiner@nyu.edu

new light scalar ϕ , with $m_\phi^2 \lesssim M_\chi \delta$. The χ can excite by emitting a ϕ with amplitude λ_- or can scatter elastically with amplitude λ_+ . We generally assume $\lambda_- \sim \lambda_+$, but this need not necessarily be the case.

The XDM dark-sector Lagrangian takes the form [3],

$$\mathcal{L}_{\text{XDM}} = \frac{1}{2} \partial_\mu \phi \partial^\mu \phi + \chi_i^\dagger \sigma_\mu \partial^\mu \chi_i - m_D \chi_1 \chi_2 - \lambda_1 \phi \chi_1^2 - \lambda_2 \phi \chi_2^2 - V(\phi) + \alpha_{\text{XDM}} \phi^2 h^2. \quad (1)$$

The last term couples the dark sector to the Standard Model via the Higgs. Defining new basis states $\chi, \chi^* = 1/\sqrt{2}(\chi_1 \mp \chi_2)$, and new couplings $\lambda_\pm = \lambda_1 \pm \lambda_2$, the Lagrangian becomes

$$\mathcal{L}_{\text{XDM}} = \frac{1}{2} \partial_\mu \phi \partial^\mu \phi + \chi^\dagger \sigma_\mu \partial^\mu \chi + \chi_*^\dagger \sigma_\mu \partial^\mu \chi_* - \frac{1}{2} (-m_D \chi^2 + m_D \chi_*^2 + \lambda_+ \phi \chi^2 + \lambda_+ \phi \chi_*^2) - \lambda_- \phi \chi \chi_* - V(\phi) + \alpha_{\text{XDM}} \phi^2 h^2. \quad (2)$$

We write $\alpha_{\text{XDM}} = m_\phi^2 K / 2v_h^2$, where K is the tuning parameter that determines the correction to the ϕ VEV and v_h is the VEV of the Higgs h ($v_h = 246$ GeV in the Standard Model): naturalness demands that $K \leq 100$ (corresponding to a 1% fine-tuning), and $\alpha_{\text{XDM}} \leq 10^{-3}$. The mass splitting δ is given by $\delta = 2\lambda_+ \langle \phi \rangle$, where $\langle \phi \rangle$ is the vacuum expectation value of the ϕ scalar. While XDM in its current form is an ad hoc extension to the Standard Model and is not supersymmetric, we expect it to be possible to realize the XDM mechanism in supersymmetric models as well.

III. EFFECT OF THE INTERMEDIATE LIGHT SCALAR ON THE NUCLEAR SCATTERING CROSS SECTION

XDM WIMPs couple to the visible sector by the mixing of the Higgs with the new light scalar ϕ , so at tree level, WIMP-nucleus scattering occurs only via t -channel ϕ -Higgs exchange. A similar scenario has been explored in [12], where the dark matter is itself a scalar that couples to the Higgs. In generic models of dark matter other scattering channels are available (such as t -channel Z exchange, and s and t -channel squark exchange in the Minimal Supersymmetric Standard Model (MSSM)), but the t -channel Higgs exchange (Fig. 1) generally provides the dominant contribution [13]. Consequently, the WIMP-nucleus scattering cross section in XDM is suppressed mostly by the mixing angle between the new light scalar and the Higgs, and the suppression factor can be calculated in a straightforward way.

The Feynman diagram for t -channel Higgs-mediated WIMP-nucleus scattering in XDM (Figure 2) is identical to Figure 1 except that the χ and χ_* WIMPs do not couple directly to the Higgs: their interaction with the Higgs is mediated by the relatively light scalar ϕ boson, which then mixes with the Higgs. Integrating out the ϕ




FIG. 1: t -channel WIMP-nucleus scattering via Higgs transfer.




FIG. 2: t -channel WIMP-nucleus scattering via ϕ -mediated Higgs transfer (XDM scenario).

propagator gives rise to an effective WIMP-Higgs interaction with amplitude $\lambda_+ 4\alpha_{\text{XDM}} v_h \langle \phi \rangle / (m_\phi^2 - t)$. The analysis is identical for elastic scattering of χ^* particles; for the inelastic de-excitation $\chi_* A \rightarrow \chi A$ (where A denotes the nucleus) the only change is that λ_+ is replaced with λ_- . (The inelastic excitation $\chi A \rightarrow \chi_* A$ is kinematically forbidden for virtually all the WIMPs in the Earth's neighborhood.) In the center-of-mass frame $t = -2|\mathbf{p}|^2(1 - \cos \theta)$, where θ is the scattering angle and the incoming WIMP has 4-momentum $\mathbf{p}$. For the energy range of interest, the momentum transfer is generally small compared to m_ϕ , so the effective coupling becomes,

$$g_{h\chi\chi} = 4\lambda_+ \alpha_{\text{XDM}} v_h \langle \phi \rangle / m_\phi^2 = \frac{\delta K}{v_h}. \quad (3)$$

(If m_ϕ is very small, of the same order as the momentum transfer, then the ϕ exchange can introduce a simple angular dependence to the cross section.)

This effective coupling is very small: allowing K to take values up to 100, we find,

$$g_{h\chi\chi} \approx 4.1 \times 10^{-4} \left(\frac{K}{100} \right) \left(\frac{\delta}{1 \text{ MeV}} \right) \left(\frac{v_h}{246 \text{ GeV}} \right)^{-1}. \quad (4)$$

Suppose we compare XDM to a generic model with $g_{h\chi\chi} = 0.1$ as a benchmark. Then we see the XDM nuclear scattering cross section (proportional to the square of the coupling) is suppressed by at least five orders of magnitude.

IV. CALCULATION OF THE XDM WIMP-NUCLEUS ELASTIC SCATTERING CROSS SECTION

We now calculate the XDM WIMP-nucleus cross section directly, for XDM coupled to the Higgs sector of the Standard Model. We will compute the effective spin-independent cross section for low-energy scattering between a WIMP and a pointlike nucleus. (To determine the actual scattering cross section this result must then be integrated over the nuclear form factor.)

Evaluating the elastic scattering amplitude $\mathcal{M}$ in the low-energy ($t \ll m_\phi \ll m_h$) nonrelativistic regime, for a nucleus with Z protons and $A - Z$ neutrons, we obtain,

$$|\mathcal{M}| = (2M)(2m_\chi) \frac{g_{h\chi\chi}}{m_h^2} (Zg_{hpp} + (A - Z)g_{hnn}), \quad (5)$$

where g_{hpp} and g_{hnn} are the couplings of the Higgs to the proton and neutron respectively, and M is the nuclear mass. Thus the cross section due to Higgs exchange is given by,

$$\sigma = \frac{\mu_{A\chi}^2}{\pi} \left(\frac{K\delta}{v_h} \right)^2 \left(\frac{Zg_{hpp} + (A - Z)g_{hnn}}{m_h^2} \right)^2, \quad (6)$$

where $\mu_{A\chi}$ is the reduced mass of the system.

The couplings of the Higgs to the proton and neutron are given by [13, 14],

$$g_{hNN} = \sum_{q=u,d,s,c,b,t} \langle N | \bar{q}q | N \rangle g_{hq} \quad (7)$$

$$\langle N | \bar{q}q | N \rangle = f_{Tq}^N m_N / m_q, \quad (8)$$

where $N = p, n$, and the nucleon parameters f_{Tq}^N are given by [15, 16, 17]

$$f_{Tu}^p = 0.023, \quad f_{Td}^p = 0.034, \quad f_{Ts}^p = 0.14, \\ f_{Tc}^p = f_{Tb}^p = f_{Tt}^p = 0.0595, \quad (9)$$

$$f_{Tu}^n = 0.019, \quad f_{Td}^n = 0.041, \quad f_{Ts}^n = 0.14,$$

$$f_{Tc}^n = f_{Tb}^n = f_{Tt}^n = 0.0592. \quad (10)$$

Note that f_{Ts}^N is not well determined [15]: values in the literature also include 0.08 [18] and 0.46 [16, 19]. We adopt the value obtained by Gasser et al [17] and used in the DarkSUSY package [14]. The nucleon parameters for the heavy quarks c, b, t are determined by those for the light quarks [15, 20], so the uncertainty in f_{Ts}^N carries over to those parameters as well. However, choosing different values for f_{Ts}^N modifies the final cross section by a factor $\sim 3 - 4$ at most.

In the Standard Model, the quark-Higgs Yukawa couplings are given by $g_{hq} = -m_q/v_h$, so the nuclear scattering cross section takes the form,

$$\sigma = \frac{K^2}{\pi} \left(\frac{\mu_{A\chi}\delta}{v_h^2 m_h^2} \right)^2 (0.3776(A - Z)m_n + 0.3755Zm_p)^2. \quad (11)$$

To compare with the current limits from direct detection experiments, we now take $A = Z = 1$ and compute the scattering cross section for a single proton. In this case $m_p = M \ll m_\chi$ so $\mu_{A\chi} \approx m_p$. The lab-frame low-energy elastic scattering cross section is then,

$$\sigma \approx 4 \times 10^{-13} \text{pb} \times \left(\frac{m_h}{100 \text{GeV}} \right)^{-4} \times \left(\frac{K}{100} \right)^2 \times \left(\frac{\delta}{1 \text{MeV}} \right)^2 \times \left(\frac{v_h}{246 \text{GeV}} \right)^{-4}. \quad (12)$$

This is a very small cross section, some five orders of magnitude below current experimental limits.

V. TWO-HIGGS-DOUBLET MODELS

If the XDM dark sector is instead coupled to a two-Higgs-doublet model (2HDM), the cross section can potentially be significantly enhanced by interactions involving the heavy Higgs. Such models constitute one of the simplest extensions of the Standard Model that allow additional CP violation [21, 22], and have been discussed extensively in the literature (see [23] for a review). We will concern ourselves here with type-II 2HDMs, where one Higgs doublet couples purely to up-type fermions and the other couples purely to down-type fermions [24]: the MSSM is one example of such a model. The effective coupling between χ and the heavy Higgs can be quite large for large $\tan\beta$ in such models, as the smaller VEV results in a weaker constraint on the ϕ -Higgs mixing α_{XDM} .

Such a model requires the Lagrangian (Eq. 2) to be modified to take couplings to the second Higgs into account. For example, let us replace the usual ϕ -Higgs mixing term with $\alpha_{\text{XDM}}\phi^2 H_u^0 H_d^0 + h.c.$, where as usual the VEVs of the neutral Higgses are defined by,

$$\langle H_u^0 \rangle = \frac{v_u}{\sqrt{2}}, \quad \langle H_d^0 \rangle = \frac{v_d}{\sqrt{2}}, \quad (13)$$

and $\tan\beta = v_u/v_d$, $v_h^2 = v_u^2 + v_d^2 \approx (246 \text{GeV})^2$. The H_u^0 - ϕ and H_d^0 - ϕ mixing terms then become, respectively, $(2\alpha_{\text{XDM}}\langle\phi\rangle v_d/\sqrt{2})\phi(H_u^0 + h.c.)$ and $(2\alpha_{\text{XDM}}\langle\phi\rangle v_u/\sqrt{2})\phi(H_d^0 + h.c.)$. The constraint on α_{XDM} now comes from the term $2(\alpha_{\text{XDM}}v_u v_d/2)\phi^2$, and can be written as,

$$\alpha_{\text{XDM}} = K \frac{m_\phi^2}{2v_u v_d}, \quad (14)$$

where as previously $K \leq 100$. Thus the Higgs- ϕ mixing terms in the Lagrangian can be written as,

$$\mathcal{L}_{\text{mixing}} = \frac{1}{\sqrt{2}} K m_\phi^2 \langle\phi\rangle \times \left(\frac{1}{v_u} H_u^0 + \frac{1}{v_d} H_d^0 \right) \phi + h.c. \quad (15)$$

We are interested in scatterings mediated by the scalar uncharged physical Higgs bosons. We now transform to

the basis of mass eigenstates, h^0 and H^0 , which as usual are related to H_u^0 and H_d^0 by a mixing angle α ,

$$\begin{aligned}\Re H_u^0 &= \frac{1}{\sqrt{2}} (\cos \alpha h^0 + \sin \alpha H^0), \\ \Re H_d^0 &= \frac{1}{\sqrt{2}} (-\sin \alpha h^0 + \cos \alpha H^0).\end{aligned}\quad (16)$$

Thus the mixing terms become,

$$\begin{aligned}\mathcal{L}_{\text{mixing}} &= Km_\phi^2 \langle \phi \rangle \times \left(\left(\frac{\cos \alpha}{v_u} - \frac{\sin \alpha}{v_d} \right) h^0 + \right. \\ &\quad \left. + \left(\frac{\sin \alpha}{v_u} + \frac{\cos \alpha}{v_d} \right) H^0 \right) \phi.\end{aligned}\quad (17)$$

The effective Higgs-WIMP interactions can then be written,

$$\begin{aligned}g_{h^0\chi\chi} &= \frac{K\delta}{2} \left(\frac{\cos \alpha}{v_u} - \frac{\sin \alpha}{v_d} \right) \\ g_{H^0\chi\chi} &= \frac{K\delta}{2} \left(\frac{\sin \alpha}{v_u} + \frac{\cos \alpha}{v_d} \right).\end{aligned}\quad (18)$$

The quark-Higgs Yukawa couplings $g_{hq\bar{q}}$ for the light CP-even Higgs are given by [15],

$$g_{hq\bar{q}} = \frac{m_q}{v_h} \times \begin{cases} -\frac{\cos \alpha}{\sin \beta} & \text{for } q = u, c, t, \\ \frac{\sin \alpha}{\cos \beta} & \text{for } q = d, s, b, \end{cases}\quad (19)$$

whereas those for the heavy CP-even Higgs are,

$$g_{Hq\bar{q}} = \frac{m_q}{v_h} \times \begin{cases} -\frac{\sin \alpha}{\sin \beta} & \text{for } q = u, c, t, \\ -\frac{\cos \alpha}{\cos \beta} & \text{for } q = d, s, b. \end{cases}\quad (20)$$

The Standard Model couplings are recovered in the decoupling limit where $\alpha \approx \beta - \pi/2$.

The effective WIMP-nucleon interaction for a proton thus becomes,

$$\begin{aligned}g_{p\chi\chi} &= \frac{K\delta m_p}{2v_h^2} \left(\frac{1}{m_{h^0}^2} \left(\frac{\cos \alpha}{\sin \beta} - \frac{\sin \alpha}{\cos \beta} \right) \times \right. \\ &\quad \times \left(0.2335 \frac{\sin \alpha}{\cos \beta} - 0.142 \frac{\cos \alpha}{\sin \beta} \right) + \\ &\quad + \frac{1}{m_{H^0}^2} \left(\frac{\sin \alpha}{\sin \beta} + \frac{\cos \alpha}{\cos \beta} \right) \times \\ &\quad \times \left(-0.142 \frac{\sin \alpha}{\sin \beta} - 0.2335 \frac{\cos \alpha}{\cos \beta} \right) \Bigg)\end{aligned}\quad (21)$$

The nuclear scattering cross section normalized to a single nucleon is, as previously,

$$\sigma = \frac{\mu_{A\chi}^2}{\pi} g_{p\chi\chi}^2.\quad (22)$$

At large $\tan \beta$, Eq. 21 can give rise to a much greater cross section than in the Standard Model case (Eq. 12), via the contribution from the heavy Higgs. In the most general two-Higgs-doublet model, the Higgs masses and




FIG. 3: Variation of the cross section with $\tan \beta$ for XDM coupled to a two-Higgs-doublet model, for $m_A = 300, 400, 500$ GeV. The thick lines correspond to cases where the light Higgs mass is held constant at 100 GeV, whereas the thin lines use the tree-level MSSM values.

mixing angles are all independent parameters: however, allowing all these parameters to vary freely can lead to conflicts with experimental constraints, and problematic flavor changing neutral current processes. To estimate the effect of increasing $\tan \beta$, we consider the tree-level MSSM relations between m_H, m_h, α and β [15],

$$\begin{aligned}m_{h,H}^2 &= \frac{1}{2} (m_A^2 + M_Z^2 \mp \\ &\quad \mp \sqrt{(m_A^2 + M_Z^2)^2 - 4M_Z^2 m_A^2 \cos^2 2\beta}),\end{aligned}\quad (23)$$

$$\frac{\sin 2\alpha}{\sin 2\beta} = -\frac{m_A^2 + m_Z^2}{m_H^2 - m_h^2}, \quad \frac{\cos 2\alpha}{\cos 2\beta} = -\frac{m_A^2 - m_Z^2}{m_H^2 - m_h^2}.\quad (24)$$

Note that these relations are only approximately correct: the Higgs masses experience significant radiative corrections.

Fig. 3 shows the variation of the spin-independent cross section as a function of $\tan \beta$, for different pseudoscalar Higgs masses m_A (see [25] for a recent analysis of constraints on m_A). We also show the cross section where the light Higgs mass is held constant at 100 GeV (but the tree-level relations are used to obtain the mixing angle α), to facilitate comparison with the Standard Model case discussed previously and to explore the effect on the nuclear scattering cross section of varying the light Higgs mass. At large $\tan \beta$ we find that varying the mass of the light Higgs has little impact, as the contribution from exchange of the heavy Higgs dominates.

which is beyond the reach of future proposed direct detection experiments. However, coupling the XDM dark sector to a two-Higgs-doublet model can give rise to interesting cross sections, provided $\tan\beta$ is large.

Models of dark matter incorporating inelastic scattering may be of interest in reconciling the results of DAMA/LIBRA with other direct detection experiments,

but XDM in its current form is unsuitable, as the mass splitting is too large for the excited state to be kinematically accessible by WIMPs in the neighborhood of the Earth. However, extensions of the model containing multiple nearly degenerate excited states might allow inelastic scattering to play a greater role.

-
- [1] G. Weidenspointner et al., *Astron. Astrophys.* **450**, 1013 (2006), astro-ph/0601673.
 - [2] G. Weidenspointner et al., arXiv:astro-ph/0702621v (2007), astro-ph/0702621v1.
 - [3] D. P. Finkbeiner and N. Weiner, *Phys. Rev.* **D76**, 083519 (2007), astro-ph/0702587.
 - [4] C. Boehm, D. Hooper, J. Silk, M. Cassé, and J. Paul, *Phys. Rev. Lett.* **92**, 101301 (2004), astro-ph/0309686.
 - [5] J. F. Beacom, N. F. Bell, and G. Bertone, *Phys. Rev. Lett.* **94**, 171301 (2005), astro-ph/0409403.
 - [6] J. F. Beacom and H. Yuksel, *Phys. Rev. Lett.* **97**, 071102 (2006), astro-ph/0512411.
 - [7] M. Pospelov and A. Ritz, *Phys. Lett.* **B651**, 208 (2007), hep-ph/0703128.
 - [8] R. Bernabei, P. Belli, F. Cappella, R. Cerulli, C. J. Dai, A. d'Angelo, H. L. He, A. Incicchitti, H. H. Kuang, J. M. Ma, et al., *ArXiv e-prints* **804** (2008), 0804.2741.
 - [9] S. Chang, G. D. Kribs, D. Tucker-Smith, and N. Weiner, *ArXiv e-prints* **807** (2008), 0807.2250.
 - [10] J. Chang et al. (ATIC) (2005), prepared for 29th International Cosmic Ray Conferences (ICRC 2005), Pune, India, 31 Aug 03 - 10 2005.
 - [11] N. Arkani-Hamed, D. Finkbeiner, T. Slatyer, and N. Weiner (2008), 0810.0713.
 - [12] V. Barger, P. Langacker, M. McCaskey, M. J. Ramsey-Musolf, and G. Shaughnessy, *Phys. Rev.* **D77**, 035005 (2008), 0706.4311.
 - [13] M. Drees and M. M. Nojiri, *Phys. Rev. D* **48**, 3483 (1993), arXiv:hep-ph/9307208.
 - [14] P. Gondolo, J. Edsjö, P. Ullio, L. Bergström, M. Schelke, and E. A. Baltz, *Journal of Cosmology and Astro-Particle Physics* **7**, 8 (2004), arXiv:astro-ph/0406204.
 - [15] G. Jungman, M. Kamionkowski, and K. Griest, *Physics Reports* **267**, 195 (1996), arXiv:hep-ph/9506380.
 - [16] H.-Y. Cheng, *Phys. Lett.* **B219**, 347 (1989).
 - [17] J. Gasser, H. Leutwyler, and M. E. Sainio, *Physics Letters B* **253**, 252 (1991).
 - [18] T. Hatsuda and T. Kunihiro, *Nuclear Physics B* **387**, 715 (1992).
 - [19] T. P. Cheng, *Phys. Rev. D* **38**, 2869 (1988).
 - [20] M. A. Shifman, A. I. Vainshtein, and V. I. Zakharov, *Physics Letters B* **78**, 443 (1978).
 - [21] T. D. Lee, *Phys. Rev. D* **8**, 1226 (1973).
 - [22] S. Weinberg, *Phys. Rev. Lett.* **37**, 657 (1976).
 - [23] E. Accomando et al. (2006), hep-ph/0608079.
 - [24] J. F. Donoghue and L.-F. Li, *Phys. Rev. D* **19**, 945 (1979).
 - [25] D. Eriksson, F. Mahmoudi, and O. Stal (2008), 0808.3551.
 - [26] D. R. Smith and N. Weiner, *Phys. Rev.* **D64**, 043502 (2001), hep-ph/0101138.
 - [27] D. Tucker-Smith and N. Weiner, *Phys. Rev.* **D72**, 063509 (2005), hep-ph/0402065.
 - [28] D. P. Finkbeiner, N. Padmanabhan, and N. Weiner, *ArXiv e-prints* **805** (2008), 0805.3531.
 - [29] N. Arkani-Hamed and N. Weiner (2008), 0810.0714.
 - [30] R. Gaitskell and J. Filippini, <http://dmttools.berkeley.edu/limitplots> (2008).